

**#MADEEASY**



# PHASES OF NERVE ACTION POTENTIAL





## Can you simplify Resting Membrane Potential (RMP)?

**Sure,**

- **Imagine a quiet room (representing neuron) with a door (representing cell membrane) almost closed but slightly open.**
- **At RMP, the neuron is at rest with a stable negative charge (around -70 MVQ) due to the distribution of ions across the cell membrane.**



# How do you open doors for nerve **Depolarization**?



***Keep in mind the previous example and imagine:***

- ***Suddenly, someone rings the doorbell (stimulus).***
- ***In response to the stimulus, the door (cell membrane) opens completely (voltage-gated sodium channels open).***
- ***Sodium ions ( $\text{Na}^+$ ) rush into the room (neuron), making the inside less negative (membrane potential now becomes less negative, moving closer to 0 mV).***



# How do you close the gate? (Repolarization Phase)



- **Now, the doorbell has stopped ringing (closing sodium channels and opening potassium channels).**
- **The room becomes quiet again (repolarization).**
- **Potassium ions ( $K^+$ ) rush out of the room (neuron), making it negative again (restoring the negative membrane potential).**





## And what happens in Hyperpolarization Phase?

***In this phase :***

- ***The quiet room is now empty, making it even quieter (slight overshoot of potassium efflux).***
- ***Excessive  $K^+$  efflux leads to a more positive membrane potential (hyperpolarization).***





# What is Refractory Period?

- **Now, the door (*cell membrane*) is completely closed and is stuck in the refractory period, no matter how loud the doorbell rings (*neuron temporarily cannot generate another action potential*).**
- **During the refractory period, no new action potential can be generated irrespective of the threshold.**

